

Compact star properties from an extended linear sigma model

Presenter:

JÁNOS TAKÁTSY

Eötvös Loránd University

Supervisor:

PÉTER KOVÁCS

Wigner RCP



MODE SNR-PWN 2019
Orléans, France

9 April 2019

Outline

1. Introduction


2. Methods

3. Results

4. Conclusion

Motivation

- ▶ QCD is unsolvable on low energies and finite densities
- ▶ Effective models are used to describe strongly interacting matter in this region
- ▶ Observations of neutron stars put constraints on dense strongly interacting matter [1,2]



Mass-radius relation from different models [1]

[1] Demorest P., et al. (2010), Nature, 467, 1081

[2] Antoniadis J., et al. (2013), Science, 340, 6131

The extended linear sigma model [3]

The Lagrangian is constructed based on linearly realized global $U(3)_L \times U(3)_R$ chiral symmetry and its explicit breaking:

- ▶ The model includes three flavours of constituent quarks
- ▶ Vector and axialvector meson nonets are included in addition to scalar and pseudoscalar mesons
- ▶ Quark–vector-meson interactions, which describe short-range repulsion are not (yet) included

Contribution from Polyakov loop potential is included:

- ▶ Polyakov loop variables constructed from gluon fields
- ▶ Modifies the Fermi-Dirac distribution function → mimics quark confinement

Applied approximations and parameter determination

Calculating the grand potential:


- ▶ One-loop fermionic contributions are considered
- ▶ Mesonic contributions are considered at tree-level
- ▶ Thermal contributions of lowest mass mesons (π, K, f_0^L) are included in the pressure

Parametrization:

- ▶ 14 unknown parameters + renormalization scale
- ▶ Determined with χ^2 fit to particle masses and decay widths
- ▶ 16 experimental masses \rightarrow fitted with curvature masses with vacuum contributions (assignment is not trivial)
- ▶ 12 experimental decay widths
- ▶ Good agreement with experimental values


Equation of state

- ▶ Compared the results with results from other models [4]
- ▶ The EoS of the eLSM is close to the interacting Walecka model at low energies
- ▶ At high energies it converges to the free quark model




[4] Scmitt A. (2010), LNP, 811

Mass-radius relation



Equation of state

- ▶ Compared the M-R diagram to free quark models with different quark masses
- ▶ $m_{\text{free}} < m_{\text{eLSM}} < m_{\mu_0}$
- ▶ The mesonic interactions have a significant effect on the M-R diagram



Conclusion

Conclusion

- ▶ The model fails to satisfy the conditions imposed by observational constraints
- ▶ Quark–meson interactions play a significant role and may push the maximum mass above $2 M_{\odot}$ by including quark–vector-meson interactions

Future work

- ▶ Making the calculations consistent by including one-loop mesonic terms
- ▶ Including interactions of fermions with vector-mesons (which is needed for short-range repulsion)
- ▶ Fitting to nuclear EoS at low energies

Thank you for your attention!



Emberi Erőforrások
Minisztériuma

SUPPORTED BY THE ÚNKP-18-2 NEW NATIONAL EXCELLENCE PROGRAM
OF THE MINISTRY OF HUMAN CAPACITIES

Lagrangian I.

\mathcal{L} constructed based on linearly realized global $U(3)_L \times U(3)_R$ symmetry and its explicit breaking

$$\begin{aligned} \mathcal{L} = & \text{Tr}[(D_\mu \Phi)^\dagger (D_\mu \Phi)] - m_0^2 \text{Tr}(\Phi^\dagger \Phi) - \lambda_1 [\text{Tr}(\Phi^\dagger \Phi)]^2 - \lambda_2 \text{Tr}(\Phi^\dagger \Phi)^2 \\ & + c_1 (\det \Phi + \det \Phi^\dagger) + \text{Tr}[H(\Phi + \Phi^\dagger)] - \frac{1}{4} \text{Tr}(L_{\mu\nu}^2 + R_{\mu\nu}^2) \\ & + \text{Tr} \left[\left(\frac{m_1^2}{2} \mathbb{I} + \Delta \right) (L_\mu^2 + R_\mu^2) \right] + i \frac{g_2}{2} (\text{Tr}\{L_{\mu\nu} [L^\mu, L^\nu]\} + \text{Tr}\{R_{\mu\nu} [R^\mu, R^\nu]\}) \\ & + \frac{h_1}{2} \text{Tr}(\Phi^\dagger \Phi) \text{Tr}(L_\mu^2 + R_\mu^2) + h_2 \text{Tr}[(L_\mu \Phi)^2 + (\Phi R_\mu)^2] + 2h_3 \text{Tr}(L_\mu \Phi R^\mu \Phi^\dagger) \\ & + \bar{\Psi} i \gamma_\mu D^\mu \Psi - g_F \bar{\Psi} (\Phi_S + i \gamma_5 \Phi_{PS}) \Psi, \end{aligned}$$

$$\begin{aligned} D^\mu \Phi &= \partial^\mu \Phi - ig_1(L^\mu \Phi - \Phi R^\mu) - ie A_e^\mu [T_3, \Phi], \\ L^{\mu\nu} &= \partial^\mu L^\nu - ie A_e^\mu [T_3, L^\nu] - \{\partial^\nu L^\mu - ie A_e^\nu [T_3, L^\mu]\}, \\ R^{\mu\nu} &= \partial^\mu R^\nu - ie A_e^\mu [T_3, R^\nu] - \{\partial^\nu R^\mu - ie A_e^\nu [T_3, R^\mu]\}, \\ D^\mu \Psi &= \partial^\mu \Psi - i G^\mu \Psi, \quad \text{with} \quad G^\mu = g_s G_a^\mu T_a. \end{aligned}$$

+ Polyakov loop potential

Lagrangian II.

the **matter** and **external** fields are

$$\Phi = \sum_{i=0}^8 (\sigma_i + i\pi_i) T_i, \quad H = \sum_{i=0}^8 h_i T_i \quad T_i : U(3) \text{ generators}$$

$$R^\mu = \sum_{i=0}^8 (\rho_i^\mu - b_i^\mu) T_i, \quad L^\mu = \sum_{i=0}^8 (\rho_i^\mu + b_i^\mu) T_i, \quad \Delta = \sum_{i=0}^8 \delta_i T_i$$

$$\Psi = (u, d, s)^T$$

non strange – strange base:

$$\xi_N = \sqrt{2/3}\xi_0 + \sqrt{1/3}\xi_8,$$

$$\xi_S = \sqrt{1/3}\xi_0 - \sqrt{2/3}\xi_8, \quad \xi \in (\sigma_i, \pi_i, \rho_i^\mu, b_i^\mu, h_i)$$

broken symmetry: non-zero condensates $\langle \sigma_{N/S} \rangle \equiv \bar{\sigma}_{N/S}$

Particle content

- Vector and Axial-vector meson nonets

$$V^\mu = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{\omega_N + \rho^0}{\sqrt{2}} & \rho^+ & K^{*+} \\ \rho^- & \frac{\omega_N - \rho^0}{\sqrt{2}} & K^{*0} \\ K^{*-} & \bar{K}^{*0} & \omega_S \end{pmatrix}^\mu \quad A^\mu = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{f_{1N} + a_1^0}{\sqrt{2}} & a_1^+ & K_1^+ \\ a_1^- & \frac{f_{1N} - a_1^0}{\sqrt{2}} & K_1^0 \\ K_1^- & \bar{K}_1^0 & f_{1S} \end{pmatrix}^\mu$$

$$\begin{aligned} \rho &\rightarrow \rho(770), K^* \rightarrow K^*(894) \\ \omega_N &\rightarrow \omega(782), \omega_S \rightarrow \phi(1020) \end{aligned}$$

$$\begin{aligned} a_1 &\rightarrow a_1(1230), K_1 \rightarrow K_1(1270) \\ f_{1N} &\rightarrow f_1(1280), f_{1S} \rightarrow f_1(1426) \end{aligned}$$

- Scalar ($\sim \bar{q}_i q_j$) and pseudoscalar ($\sim \bar{q}_i \gamma_5 q_j$) meson nonets

$$\Phi_S = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{\sigma_N + a_0^0}{\sqrt{2}} & a_0^+ & K_0^{*+} \\ a_0^- & \frac{\sigma_N - a_0^0}{\sqrt{2}} & K_0^{*0} \\ K_0^{*-} & \bar{K}_0^{*0} & \sigma_S \end{pmatrix} \quad \Phi_{PS} = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{\eta_N + \pi^0}{\sqrt{2}} & \pi^+ & K^+ \\ \pi^- & \frac{\eta_N - \pi^0}{\sqrt{2}} & K^0 \\ K^- & \bar{K}^0 & \eta_S \end{pmatrix}$$

unknown assignment
mixing in the $\sigma_N - \sigma_S$ sector

$\pi \rightarrow \pi(138), K \rightarrow K(495)$
mixing: $\eta_N, \eta_S \rightarrow \eta(548), \eta'(958)$

Spontaneous symmetry breaking: $\sigma_{N/S}$ acquire nonzero expectation values $\phi_{N/S}$
fields shifted by their expectation value: $\sigma_{N/S} \rightarrow \sigma_{N/S} + \phi_{N/S}$

Polyakov loops in Polyakov gauge

Polyakov loop variables: $\Phi(\vec{x}) = \frac{\text{Tr}_c L(\vec{x})}{N_c}$ and $\bar{\Phi}(\vec{x}) = \frac{\text{Tr}_c \bar{L}(\vec{x})}{N_c}$ with
 $L(x) = \mathcal{P} \exp \left[i \int_0^\beta d\tau G_4(\vec{x}, \tau) \right]$

→ signals center symmetry (\mathbb{Z}_3) breaking at the deconfinement transition

low T : confined phase, $\langle \Phi(\vec{x}) \rangle, \langle \bar{\Phi}(\vec{x}) \rangle = 0$

high T : deconfined phase, $\langle \Phi(\vec{x}) \rangle, \langle \bar{\Phi}(\vec{x}) \rangle \neq 0$

- ▶ Polyakov gauge: $G_4(\vec{x}, \tau) = G_4(\vec{x})$, plus gauge rotation to diagonal form in color space
- ▶ further simplification: \vec{x} -independence

$$\hookrightarrow L = e^{i\beta G_4} = \text{diag}(a, b, c) \left(\begin{smallmatrix} ! \\ \in SU(3)^{\text{color}} \end{smallmatrix} \right); \quad a, b, c \in \mathbb{Z}$$

→ use this to calculate partition function of free quarks

Form of the potential

I.) Simple polynomial potential invariant under \mathbb{Z}_3 and charge conjugation: R.D.Pisarski, PRD 62, 111501

$$\frac{\mathcal{U}_{\text{poly}}^{\text{YM}}(\Phi, \bar{\Phi})}{T^4} = -\frac{b_2(T)}{2}\bar{\Phi}\Phi - \frac{b_3}{6}(\Phi^3 + \bar{\Phi}^3) + \frac{b_4}{4}(\bar{\Phi}\Phi)^2$$

with $b_2(T) = a_0 + a_1 \frac{T_0}{T} + a_2 \frac{T_0^2}{T^2} + a_3 \frac{T_0^3}{T^3}$

II.) Logarithmic potential coming from the $SU(3)$ Haar measure of group integration K. Fukushima, Phys. Lett. **B591**, 277 (2004)

$$\frac{\mathcal{U}_{\log}^{\text{YM}}(\Phi, \bar{\Phi})}{T^4} = -\frac{1}{2}a(T)\Phi\bar{\Phi} + b(T) \ln \left[1 - 6\Phi\bar{\Phi} + 4(\Phi^3 + \bar{\Phi}^3) - 3(\Phi\bar{\Phi})^2 \right]$$

with $a(T) = a_0 + a_1 \frac{T_0}{T} + a_2 \frac{T_0^2}{T^2}, \quad b(T) = b_3 \frac{T_0^3}{T^3}$

$\mathcal{U}^{\text{YM}}(\Phi, \bar{\Phi})$ models the free energy of a pure gauge theory

Determination of the parameters

14 unknown parameters ($m_0, \lambda_1, \lambda_2, c_1, m_1, g_1, g_2, h_1, h_2, h_3, \delta_S, \Phi_N, \Phi_S, g_F$) → determined by the min. of χ^2 :

$$\chi^2(x_1, \dots, x_N) = \sum_{i=1}^M \left[\frac{Q_i(x_1, \dots, x_N) - Q_i^{\text{exp}}}{\delta Q_i} \right]^2,$$

$(x_1, \dots, x_N) = (m_0, \lambda_1, \lambda_2, \dots)$, $Q_i(x_1, \dots, x_N)$ → from the model, Q_i^{exp} → PDG value, $\delta Q_i = \max\{5\%, \text{PDG value}\}$
multiparametric minimalization → MINUIT

- ▶ Curvature masses → 16 physical quantities:
 $m_u/d, m_s, m_\pi, m_\eta, m_{\eta'}, m_K, m_\rho, m_\Phi, m_{K^\star}, m_{a_1}, m_{f_1^H}, m_{K_1},$
 $m_{a_0}, m_{K_s}, m_{f_0^L}, m_{f_0^H}$
- ▶ Decay widths → 12 physical quantities:
 $\Gamma_{\rho \rightarrow \pi\pi}, \Gamma_{\Phi \rightarrow KK}, \Gamma_{K^\star \rightarrow K\pi}, \Gamma_{a_1 \rightarrow \pi\gamma}, \Gamma_{a_1 \rightarrow \rho\pi}, \Gamma_{f_1 \rightarrow KK^\star}, \Gamma_{a_0}, \Gamma_{K_S \rightarrow K\pi},$
 $\Gamma_{f_0^L \rightarrow \pi\pi}, \Gamma_{f_0^L \rightarrow KK}, \Gamma_{f_0^H \rightarrow \pi\pi}, \Gamma_{f_0^H \rightarrow KK}$

Result of the parametrization

- 40 possible assignments of scalar mesons to the scalar nonet states
- 3 values of M_0 are used \implies 120 cases to investigate
for each case $5 \cdot 10^4 - 10^5$ configurations are used for the χ^2 minimization
- lowest χ^2 obtained for $M_0 = 0.3$ GeV $\chi^2 = 18.57$ and $\chi^2_{\text{red}} \equiv \frac{\chi^2}{N_{\text{dof}}} = 1.16$
assignment: $a_0^{\bar{q}q} \rightarrow a_0(980)$, $K_0^{*,\bar{q}q} \rightarrow K_0^*(800)$, $f_0^{L,\bar{q}q} \rightarrow f_0(500)$, $f_0^{H,\bar{q}q} \rightarrow f_0(980)$
- problems: $m_{a_0} < m_{K_0^*}$, $m_{f_0^{H/L}}$ too light
- by minimizing also for M_0 we obtain using $\mathcal{U}_{\log}^{\text{YM}}(\Phi, \bar{\Phi})$ with $T_0 = 182$ MeV:

| Parameter | Value | Parameter | Value |
|------------------------|-------------|-------------|---------|
| ϕ_N [GeV] | 0.1411 | g_1 | 5.6156 |
| ϕ_S [GeV] | 0.1416 | g_2 | 3.0467 |
| m_0^2 [GeV 2] | $2.3925E-4$ | h_1 | 27.4617 |
| m_1^2 [GeV 2] | $6.3298E-8$ | h_2 | 4.2281 |
| λ_1 | -1.6738 | h_3 | 5.9839 |
| λ_2 | 23.5078 | g_F | 4.5708 |
| c_1 [GeV] | 1.3086 | M_0 [GeV] | 0.3511 |
| δ_S [GeV 2] | 0.1133 | | |